

APPENDIX 1 - vaccinations

Age due	Vaccine Given:
Eight weeks old	Diphtheria, tetanus, pertussis, polio, Haemophilus influenza type b (Hib) and Hepatitis B (DTaP/IPV/Hib/HepB) Meningococcal B (MenB) Rotavirus
Twelve weeks old	Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B (DTaP/IPV/Hib/HepB) Rotavirus Pneumococcal conjugate vaccine (PCV13) One injection One oral
Sixteen weeks old	Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B (DTaP/IPV/Hib/HepB) Meningococcal B (MenB)
One year old (on or after the child's first birthday)	Hib/MenC Pneumococcal conjugate vaccine (PCV13) Meningococcal B (MenB) Measles, mumps and rubella (MMR)
Primary school age children (school years reception to six)	Live attenuated influenza vaccine (LAIV)
Three years four months old or soon after	Diphtheria, tetanus, pertussis and polio (dTaP/IPV) Measles, mumps and rubella (MMR)
Twelve to Thirteen years	Human papillomavirus (HPV)
Fourteen years old (school year 9)	Tetanus, diphtheria and polio (Td/IPV) Meningococcal ACWY conjugate (MenACWY)
65 years old	Pneumococcal polysaccharide vaccine (PPV)
65 years of age and older	Inactivated influenza vaccine
70 years old	Shingles

Pregnant women should be vaccinated against influenza (during flu season) and pertussis