

NATIONAL GENERAL PRACTICE PROFILES

PROFILE FOR

KENTON MEDICAL CENTRE

7 NORTHWICK AVENUE, KENTON, HARROW, MIDDLESEX, HA3 0AA






These profiles are designed to support GPs, clinical commissioning groups (CCGs) and PCTs to ensure that they are providing and commissioning effective and appropriate healthcare services for their local population. Using a variety of graphical displays such as spine charts and population pyramids, the tool presents a range of practice-level indicators drawn from the latest available data, including:

- local demography;
- Quality and Outcomes Framework domains;
- disease prevalence estimates;
- admission rates; and
- patient satisfaction.

In addition to displaying individual practice profiles, the web tool allows you to view summary profiles for PCTs and CCGs. Each practice can be compared with its PCT, CCG and with England, and also with the practices in the same deprivation decile and 'peer group' (although PDF generation is limited to PCT, CCG and England comparisons). The profiles do not provide an exhaustive list of primary care indicators, but they do allow a consistent approach to comparing and benchmarking across England. More indicators will be incorporated as the tool is developed further.

The profiles have been designed as a web tool and the full functionality is only available via the web version. For more information consult the User guide and FAQs via the Supporting documents tab on the web pages.

The development of this tool has been led by erpho, the East of England Public Health Observatory, on behalf of the network of PHOs in England. For further information contact: feedback@erpho.org.uk



www.apho.org.uk/PracProf

How to read the indicator spine charts

The light grey bar shows the range of values found in England. The dark grey sections mark out the range within which the middle half of the observed values lie (25th to 75th percentile).

The red line shows where the England average is. The position of the circle shows the practice value, a diamond the PCT value, a triangle the cluster value, in relation to this scale.

The corresponding numbers can be found

in the cells next to the chart.

If significance has been calculated for the indicator, then it is determined by whether the practice value is significantly higher or lower than the England average using 95% confidence intervals.

A significant difference is marked in blue, non-significance in yellow.

A significant difference is marked in blue, non-significance in yellow.

- No significant difference from England average
- Significantly different from England average
- Significance not calculated

- Practice
- PCT
- Clinical Commissioning Group



Practice Summary

| Indicator | Period | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|---|---------|----------------|-----------|-----------------|----------------|---------------|-----------------|
| % aged 0 to 4 years | 2011 | 4.5% | 6.7% | 6.0% | 0.0% | | 16.5% |
| % aged 5 to 14 years | 2011 | 8.0% | 11.0% | 11.2% | 0.0% | | 28.0% |
| % aged under 15 years | 2011 | 12.4% | 17.7% | 17.1% | 0.0% | | 43.2% |
| % aged 65+ years | 2011 | 10.7% | 9.9% | 15.9% | 0.0% | | 45.8% |
| % aged 75+ years | 2011 | 4.7% | 4.4% | 7.5% | 0.0% | | 26.6% |
| % aged 85+ years | 2011 | 1.4% | 1.1% | 2.2% | 0.0% | | 14.3% |
| Deprivation score (IMD) | 2011 | 16.3 | 30.5 | 21.5 | 2.9 | | 68.5 |
| IDACI (Income Deprivation Affecting Children) | 2011 | 0.202 | 0.393 | 0.218 | 0.011 | | 0.684 |
| IDAOP (Income Deprivation Affecting Older People) | 2011 | 0.215 | 0.305 | 0.181 | 0.038 | | 0.802 |
| % satisfied with phone access | 2010/11 | 76.1% | 69.4% | 74.5% | 17.3% | | 100% |
| % able to see a doctor within 2 days | 2010/11 | 81.4% | 75.3% | 79.7% | 24.8% | | 100% |
| % able to book appointment >= 2d ahead | 2010/11 | 75.0% | 64.2% | 69.7% | 0.0% | | 100% |
| % satisfied with opening hours | 2010/11 | 65.8% | 73.0% | 78.3% | 43.9% | | 100% |
| % able to see preferred GP | 2010/11 | 39.3% | 61.4% | 69.8% | 11.8% | | 100% |
| % would recommend practice | 2010/11 | 66.7% | 72.0% | 83.5% | 34.0% | | 100% |
| Total QOF points | 2010/11 | 96.9% | 92.2% | 94.7% | 39.2% | | 100% |

Modelled Disease Prevalence

| Indicator | Period | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|---|---------|----------------|-----------|-----------------|----------------|---------------|-----------------|
| Estimated prevalence of CVD | 2011 | 7.0% | 7.8% | 9.5% | 2.6% | | 19.4% |
| Estimated prevalence of CHD | 2011 | 3.3% | 3.9% | 4.7% | 0.0% | | 12.9% |
| Ratio of recorded vs expected CHD prevalence | 2010/11 | 0.769 | 0.540 | 0.723 | 0.00 | | 1.87 |
| Estimated prevalence of COPD | 2011 | 2.7% | 3.2% | 2.9% | 0.62% | | 6.47% |
| Ratio of recorded vs expected COPD prevalence | 2010/11 | 0.102 | 0.220 | 0.559 | 0.00 | | 2.54 |
| Estimated prevalence of hypertension | 2011 | 21.5% | 22.8% | 24.9% | 4.2% | | 44.7% |
| Ratio of recorded vs expected hypertension prevalence | 2010/11 | 0.513 | 0.506 | 0.543 | 0.01 | | 1.14 |
| Estimated prevalence of stroke | 2011 | 1.3% | 1.7% | 2.1% | 0.16% | | 5.44% |
| Ratio of recorded vs expected stroke prevalence | 2010/11 | 0.610 | 0.588 | 0.827 | 0.00 | | 2.15 |

CVD - Coronary heart disease

| Indicator | Period | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|--|---------|----------------|-----------|-----------------|----------------|---------------|-----------------|
| CHD: QOF prevalence (all ages) | 2010/11 | 2.5% | 2.1% | 3.4% | 0.00% | | 9.89% |
| Estimated prevalence of CHD | 2011 | 3.3% | 3.9% | 4.7% | 0.0% | | 12.9% |
| Ratio of recorded vs expected CHD prevalence | 2010/11 | 0.769 | 0.540 | 0.723 | 0.00 | | 1.87 |
| Heart failure w LVD: QOF prevalence | 2010/11 | 0.1% | 0.2% | 0.4% | 0.00% | | 2.63% |
| Exception rate for CHD indicators | 2010/11 | 7.5% | 6.9% | 7.5% | 0.0% | | 100% |
| CHD Emergency Admission Rate (per 1000) | 2009/10 | 3.00 | 2.30 | 2.50 | 0.0 | | 13.9 |
| CHD Elective Admission Rate (per 1000) | 2009/10 | | 1.80 | 1.80 | 0.00 | | 10.00 |
| CHD 2: Angina referred for exercise testing &/or assessm. | 2010/11 | 92.3% | 93.2% | 94.6% | 0.0% | | 100% |
| CHD 5: Record of BP in the previous 15mths | 2010/11 | 100% | 97.3% | 97.8% | 77.6% | | 100% |
| CHD 6: Last BP reading in last 15mths is <=150/90 | 2010/11 | 90.5% | 89.5% | 90.2% | 46.5% | | 100% |
| CHD 7: Record of total cholesterol in last 15mths | 2010/11 | 93.3% | 92.0% | 93.7% | 50.0% | | 100% |
| CHD 8: Last total cholesterol is <=5mmol/l | 2010/11 | 84.0% | 81.3% | 82.1% | 40.0% | | 100% |
| CHD 9: Record that aspirin, APT or ACT is taken | 2010/11 | 98.6% | 94.3% | 93.5% | 62.5% | | 100% |
| CHD 10: Currently treated with beta blocker | 2010/11 | 88.5% | 74.0% | 73.5% | 28.6% | | 100% |
| CHD 11: History of MI: treated with ACE-I | 2010/11 | 100% | 89.2% | 88.8% | 40.0% | | 100% |
| CHD 12: CHD patients given flu immunisation 1 Sep - 31 Mar | 2010/11 | 98.4% | 92.5% | 92.4% | 40.0% | | 100% |

CVD - Stroke and TIA

| Indicator | Period | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|--|---------|----------------|-----------|-----------------|----------------|---------------|-----------------|
| Stroke: QOF prevalence (all ages) | 2010/11 | 0.8% | 1.0% | 1.7% | 0.00% | | 5.90% |
| Estimated prevalence of stroke | 2011 | 1.3% | 1.7% | 2.1% | 0.16% | | 5.44% |
| Ratio of recorded vs expected stroke prevalence | 2010/11 | 0.610 | 0.588 | 0.827 | 0.00 | | 2.15 |
| Exception rate for stroke indicators | 2010/11 | 8.3% | 6.9% | 7.1% | 0.0% | | 100% |
| Stroke 13: New patients referred for further investigation | 2010/11 | 75.0% | 84.8% | 89.8% | 0.0% | | 100% |
| Stroke 5: BP recorded in last 15mths | 2010/11 | 91.7% | 96.3% | 96.8% | 73.1% | | 100% |
| Stroke 6: Last BP reading is 150/90 or less | 2010/11 | 77.3% | 87.0% | 88.6% | 0.0% | | 100% |
| Stroke 7: Total cholesterol recorded in last 15mths | 2010/11 | 100% | 90.0% | 91.5% | 48.4% | | 100% |
| Stroke 8: Last measured total cholesterol <=5mmol/l | 2010/11 | 77.3% | 75.3% | 77.3% | 0.0% | | 100% |
| Stroke 12: Record of aspirin, APT or ACT taken | 2010/11 | 93.8% | 94.5% | 93.7% | 55.0% | | 100% |
| Stroke 10: Influenza immunisation given 1 Sep-31 Mar | 2010/11 | 100% | 90.7% | 89.6% | 0.0% | | 100% |

CVD - Heart failure and atrial fibrillation

| Indicator | Period | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|--|---------|----------------|-----------|-----------------|----------------|---------------|-----------------|
| Heart failure: QOF prevalence (all ages) | 2010/11 | 0.2% | 0.5% | 0.7% | 0.00% | | 3.88% |
| Heart failure w LVD: QOF prevalence | 2010/11 | 0.1% | 0.2% | 0.4% | 0.00% | | 2.63% |
| Exception rate for heart failure indicators | 2010/11 | 20.0% | 14.5% | 14.8% | 0.0% | | 100% |
| Atrial fibrillation: QOF prevalence | 2010/11 | 0.6% | 0.6% | 1.4% | 0.00% | | 5.19% |
| Exception rate for atrial fibrillation indicators | 2010/11 | 7.1% | 4.4% | 3.8% | 0.0% | | 100% |
| HF 2: Diagnosis conf. by ECG/specialist assessm. | 2010/11 | 50.0% | 93.6% | 95.8% | 0.0% | | 100% |
| HF 3: HF w LVD: treated with ACE-I or ARB | 2010/11 | 100% | 89.8% | 89.5% | 0.0% | | 100% |
| HF 4: Heart failure w LVD: treatment w ACE inh. or ARB, and beta-blocker | 2010/11 | 100% | 82.9% | 82.1% | 0.0% | | 100% |
| AF 4: diagnosed with ECG or by specialist | 2010/11 | 100% | 96.2% | 96.1% | 0.0% | | 100% |
| AF 3: treated w anti- coag./platelet therapy | 2010/11 | 94.1% | 94.5% | 93.6% | 0.0% | | 100% |

CVD - Risk factors for CVD

| Indicator | Period | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|---|---------|----------------|-----------|-----------------|----------------|---------------|-----------------|
| Hypertension: QOF prevalence (all ages) | 2010/11 | 11.0% | 11.5% | 13.5% | 0.1% | | 37.4% |
| Estimated prevalence of hypertension | 2011 | 21.5% | 22.8% | 24.9% | 4.2% | | 44.7% |
| Ratio of recorded vs expected hypertension prevalence | 2010/11 | 0.513 | 0.506 | 0.543 | 0.01 | | 1.14 |
| Exception rate for hypertension indicators | 2010/11 | 0.9% | 2.5% | 2.5% | 0.0% | | 22.5% |
| Exception rate for smoking indicators | 2010/11 | 0.8% | 0.8% | 0.7% | 0.00% | | 8.75% |
| Obesity: QOF prevalence (16+) | 2010/11 | 6.8% | 9.3% | 10.5% | 0.4% | | 31.0% |
| BP 4: Record of BP in last 9mths | 2010/11 | 92.0% | 90.8% | 91.6% | 66.8% | | 100% |
| BP 5: Last (9mths) blood pressure <=150/90 | 2010/11 | 75.4% | 78.3% | 79.3% | 35.1% | | 100% |
| PP 1: CV risk assessment for new hypertension cases | 2010/11 | 100% | 76.9% | 80.2% | 0.0% | | 100% |
| PP 2: life style advice for new hypertension cases | 2010/11 | 94.4% | 81.8% | 81.9% | 0.0% | | 100% |
| Dep 1: Depression case finding in CHD and/or diabetes patients | 2010/11 | 97.6% | 90.2% | 88.8% | 0.2% | | 100% |
| Smoking 3: status recorded in last 15mths (certain conditions) | 2010/11 | 99.6% | 95.8% | 95.4% | 66.7% | | 100% |
| Smoking 4: cessation advice/referral offered (certain conditions) | 2010/11 | 97.4% | 92.6% | 92.9% | 47.6% | | 100% |

Diabetes

| Indicator | Period | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|---|---------|----------------|-----------|-----------------|----------------|---------------|-----------------|
| Diabetes: QOF prevalence (17+) | 2010/11 | 6.7% | 6.9% | 5.5% | 0.0% | | 15.2% |
| Exception rate for diabetes indicators | 2010/11 | 5.6% | 6.1% | 6.5% | 0.3% | | 100% |
| Hypertension: QOF prevalence (all ages) | 2010/11 | 11.0% | 11.5% | 13.5% | 0.1% | | 37.4% |
| Exception rate for hypertension indicators | 2010/11 | 0.9% | 2.5% | 2.5% | 0.0% | | 22.5% |
| Exception rate for smoking indicators | 2010/11 | 0.8% | 0.8% | 0.7% | 0.00% | | 8.75% |
| Obesity: QOF prevalence (16+) | 2010/11 | 6.8% | 9.3% | 10.5% | 0.4% | | 31.0% |
| Diabetes admission rate (per 1000) | 2009/10 | 0.000 | 0.900 | 1.10 | 0.0 | | 10.9 |
| Ratio of recorded vs expected diabetes prevalence | 2008/09 | 1.44 | 1.38 | 0.883 | 0.19 | | 5.79 |
| DM 2: Record of BMI in the last 15mths | 2010/11 | 97.0% | 94.1% | 94.8% | 52.0% | | 100% |
| DM 5: Record of HbA1c/equivalent last 15mths | 2010/11 | 95.8% | 95.4% | 97.5% | 63.6% | | 100% |
| DM 23: Last HbA1c is <=7 in last 15mths | 2010/11 | 47.4% | 47.9% | 54.2% | 0.0% | | 99.4% |
| DM 24: Last HbA1c is <=8 in last 15mths | 2010/11 | 79.6% | 73.7% | 78.0% | 2.7% | | 100% |
| DM 25: Last HbA1c is <=9 in last 15mths | 2010/11 | 90.0% | 85.0% | 88.4% | 2.7% | | 100% |
| DM 21: Retinal screening in last 15mths | 2010/11 | 90.1% | 88.2% | 91.6% | 26.1% | | 100% |
| DM 9: Record of peripheral pulses last 15mths | 2010/11 | 92.5% | 91.8% | 91.6% | 9.6% | | 100% |
| DM 10: Record of neuropathy test last 15mths | 2010/11 | 92.5% | 91.6% | 91.4% | 22.8% | | 100% |
| DM 11: Record of BP in last 15mths | 2010/11 | 98.8% | 97.6% | 98.4% | 78.8% | | 100% |
| DM 12: Last BP is <=145/85 | 2010/11 | 80.6% | 81.4% | 81.2% | 34.6% | | 100% |
| BP 4: Record of BP in last 9mths | 2010/11 | 92.0% | 90.8% | 91.6% | 66.8% | | 100% |
| BP 5: Last (9mths) blood pressure <=150/90 | 2010/11 | 75.4% | 78.3% | 79.3% | 35.1% | | 100% |
| DM 13: Record of micro-albuminuria test last 15mths | 2010/11 | 93.1% | 88.3% | 89.0% | 0.0% | | 100% |
| DM 22: eGRF or serum creatinin testing in last 15mths | 2010/11 | 94.6% | 95.4% | 97.2% | 56.3% | | 100% |
| DM 15: Proteinuria/micro-album. treated w inhibitors | 2010/11 | 100% | 89.1% | 88.2% | 0.0% | | 100% |
| DM 16: Record of total cholesterol last 15mths | 2010/11 | 95.8% | 94.7% | 96.1% | 67.3% | | 100% |
| DM 17: Measured total chol (last 15mths) <=5mmol/l | 2010/11 | 80.7% | 80.4% | 82.9% | 46.0% | | 100% |
| DM 18: Influenza immunisation given 1 Sep - 31 Mar | 2010/11 | 94.9% | 89.9% | 90.7% | 47.1% | | 100% |
| Dep 1: Depression case finding in CHD and/or diabetes patients | 2010/11 | 97.6% | 90.2% | 88.8% | 0.2% | | 100% |
| Smoking 3: status recorded in last 15mths (certain conditions) | 2010/11 | 99.6% | 95.8% | 95.4% | 66.7% | | 100% |
| Smoking 4: cessation advice/referral offered (certain conditions) | 2010/11 | 97.4% | 92.6% | 92.9% | 47.6% | | 100% |

Mental Health

| Indicator | Period | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|--|---------|----------------|-----------|-----------------|----------------|---------------|-----------------|
| Psychoses: QOF prevalence (all ages) | 2010/11 | 0.7% | 1.0% | 0.8% | 0.0% | | 23.8% |
| Exception rate for patients with psychoses indicators | 2010/11 | 0.0% | 6.1% | 10.4% | 0.0% | | 81.5% |
| Dementia: QOF prevalence (all ages) | 2010/11 | 0.0% | 0.3% | 0.5% | 0.0% | | 10.5% |
| Exception rate for dementia indicators | 2010/11 | 0.0% | 10.2% | 7.4% | 0.0% | | 100% |
| Depression: QOF prevalence (18+) | 2010/11 | 2.9% | 5.9% | 11.2% | 0.0% | | 46.1% |
| Exception rate for depression indicators | 2010/11 | 3.2% | 4.3% | 5.9% | 0.0% | | 52.9% |
| MH 9: full review in last 15mths | 2010/11 | 100% | 91.6% | 92.6% | 0.0% | | 100% |
| MH 4: Li-therapy: record of s.creat. & TSH last 15mths | 2010/11 | 100% | 95.7% | 98.1% | 0.0% | | 100% |
| MH 5: Lithium therapy: record of Li-levels last 6mths | 2010/11 | 100% | 85.3% | 92.2% | 0.0% | | 100% |
| MH 6: with comprehensive care plan | 2010/11 | 100% | 92.0% | 89.4% | 0.0% | | 100% |
| MH 7: Pychoses patients who DNA review: followed up < 14 d | 2010/11 | % | 81.9% | 94.6% | 0.0% | | 100% |
| Dem 2: Dementia care has been reviewed last 15mths | 2010/11 | 100% | 83.2% | 79.2% | 0.0% | | 100% |
| Dep 1: Depression case finding in CHD and/or diabetes patients | 2010/11 | 97.6% | 90.2% | 88.8% | 0.2% | | 100% |
| Dep 2: Depression cases with severity assessment | 2010/11 | 100% | 86.0% | 92.1% | 0.0% | | 100% |
| Dep 3: second severity assessment for new depression cases | 2010/11 | 100% | 69.1% | 74.5% | 0.0% | | 100% |

Respiratory Disease

| Indicator | Period | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|---|---------|----------------|-----------|-----------------|----------------|---------------|-----------------|
| COPD: QOF prevalence (all ages) | 2010/11 | 0.3% | 0.7% | 1.6% | 0.00% | | 7.81% |
| Estimated prevalence of COPD | 2011 | 2.7% | 3.2% | 2.9% | 0.62% | | 6.47% |
| Ratio of recorded vs expected COPD prevalence | 2010/11 | 0.102 | 0.220 | 0.559 | 0.00 | | 2.54 |
| Exception rate for COPD indicators | 2010/11 | 0.0% | 11.3% | 12.4% | 0.0% | | 100% |
| COPD admission rate (per 1000) | 2009/10 | | 0.900 | 2.00 | 0.0 | | 17.2 |
| Asthma: QOF prevalence (all ages) | 2010/11 | 3.4% | 4.4% | 5.9% | 0.0% | | 12.4% |
| Exception rate for asthma indicators | 2010/11 | 1.7% | 1.9% | 5.5% | 0.0% | | 100% |
| Exception rate for smoking indicators | 2010/11 | 0.8% | 0.8% | 0.7% | 0.00% | | 8.75% |
| COPD 12: Diagnosis conf. by spirometry in last 15mths | 2010/11 | 100% | 87.7% | 89.7% | 0.0% | | 100% |
| COPD 10: Record of FeV1 in last 15mths | 2010/11 | 100% | 87.1% | 88.9% | 0.0% | | 100% |
| COPD 13: assessed using MRC dyspnoea score last 15mths | 2010/11 | 100% | 90.2% | 91.6% | 0.0% | | 100% |
| COPD 8: Influenza immunisation given 1 Sep - 31 Mar | 2010/11 | 100% | 92.8% | 93.2% | 50.0% | | 100% |
| Asthma 8: with measures of variability/reversibility (8+) | 2010/11 | 100% | 89.5% | 87.7% | 0.0% | | 100% |
| Asthma 3: smoking recorded in last 15 mths (14-19y w asthma) | 2010/11 | 100% | 90.5% | 89.2% | 20.0% | | 100% |
| Asthma 6: review in the last 15mths | 2010/11 | 80.0% | 80.6% | 78.6% | 32.4% | | 100% |
| Smoking 3: status recorded in last 15mths (certain conditions) | 2010/11 | 99.6% | 95.8% | 95.4% | 66.7% | | 100% |
| Smoking 4: cessation advice/referral offered (certain conditions) | 2010/11 | 97.4% | 92.6% | 92.9% | 47.6% | | 100% |





























Chronic Kidney Disease

| Indicator | Period | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|--|---------|----------------|-----------|-----------------|----------------|---------------|-----------------|
| CKD: QOF prevalence (18+) | 2010/11 | 0.6% | 2.0% | 4.3% | 0.0% | | 21.5% |
| Exception rate for CKD indicators | 2010/11 | 1.9% | 3.9% | 4.0% | 0.0% | | 100% |
| CKD 2: Record of BP in last 15mths | 2010/11 | 93.8% | 97.6% | 97.5% | 76.9% | | 100% |
| CKD 3: Last BP reading measured in last 15mths is <=140/85 | 2010/11 | 87.5% | 75.6% | 74.2% | 0.0% | | 100% |
| CKD 5: Hypertension treated with ACE inhibitor/ARB | 2010/11 | 100% | 91.2% | 90.5% | 0.0% | | 100% |
| CKD 6: Urine albumin : creatinine ratio test last 15mths | 2010/11 | 81.3% | 82.6% | 82.2% | 0.0% | | 100% |

Other Conditions

| Indicator | Period | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|---|---------|----------------|-----------|-----------------|----------------|---------------|-----------------|
| Cancer: QOF prevalence (all ages) | 2010/11 | 1.0% | 1.0% | 1.6% | 0.00% | | 4.51% |
| Exception rate for cancer indicators | 2010/11 | 0.0% | 1.4% | 1.6% | 0.0% | | 100% |
| Cancer admission rate (per 1000) | 2009/10 | 18.0 | 16.7 | 28.1 | 1.3 | | 105.2 |
| Epilepsy: QOF prevalence (18+) | 2010/11 | 0.1% | 0.5% | 0.8% | 0.00% | | 3.76% |
| Exception rate for epilepsy indicators | 2010/11 | 0.0% | 10.0% | 8.1% | 0.0% | | 66.7% |
| Learning disability: QOF prevalence (18+) | 2010/11 | 0.2% | 0.3% | 0.4% | 0.00% | | 5.86% |
| Hypothyroidism: QOF prevalence (all ages) | 2010/11 | 2.7% | 1.7% | 3.0% | 0.00% | | 7.88% |
| Exception rate for hypothyroidism indicators | 2010/11 | 0.0% | 0.5% | 0.5% | 0.0% | | 100% |
| Palliative/supportive care: QOF prevalence (all ages) | 2010/11 | 0.0% | 0.1% | 0.2% | 0.00% | | 2.76% |
| Cancer 3: review within 6mths of diagnosis | 2010/11 | 100% | 92.5% | 93.7% | 0.0% | | 100% |
| Epilepsy 6: Record of seizure frequency | 2010/11 | 100% | 96.8% | 95.6% | 22.0% | | 100% |
| Epilepsy 7: Medication review w patient/carer last 15mths | 2010/11 | 100% | 97.0% | 95.2% | 27.5% | | 100% |
| Epilepsy 8: Seizure free for last 12mths | 2010/11 | 100% | 70.6% | 73.9% | 0.0% | | 100% |
| Thyroid 2: function test recorded last 15mths | 2010/11 | 97.5% | 95.0% | 95.9% | 71.1% | | 100% |

Secondary Care Use

| Indicator | Period | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|--|---------|----------------|-----------|-----------------|----------------|---|-----------------|
| All GP referrals (1st attendance) (per 1000) | 2009/10 | 153.7 | 172.5 | 186.8 | 34.0 |  | 615.0 |
| All attendances (per 1000) | 2009/10 | 661.2 | 705.4 | 739.9 | 105 |  | 1582 |
| Dermatology, GP referrals (1st attendance) (per 1000) | 2009/10 | 4.90 | 10.1 | 13.7 | 0.0 |  | 71.8 |
| General Medicine, GP referrals (1st attendance) (per 1000) | 2009/10 | 6.20 | 7.00 | 10.9 | 0.0 |  | 81.3 |
| General Medicine - return ratio | 2009/10 | 3.60 | 4.10 | 2.70 | 0.5 |  | 31.0 |
| General Surgery, GP referrals (1st attendance) (per 1000) | 2009/10 | 13.1 | 18.4 | 23.1 | 1.9 |  | 70.8 |
| General Surgery - return ratio | 2009/10 | 1.50 | 1.40 | 1.50 | 0.30 |  | 5.10 |
| Gynaecology, GP referrals (1st attendance) (per 1000) | 2009/10 | 19.3 | 17.9 | 17.1 | 2.2 |  | 56.4 |
| Gynaecology - return ratio | 2009/10 | 1.40 | 1.50 | 1.30 | 0.20 |  | 8.50 |
| Orthopaedics, GP referrals (1st attendance) (per 1000) | 2009/10 | 16.1 | 17.1 | 20.2 | 0.0 |  | 81.4 |
| Orthopaedics - return ratio | 2009/10 | 2.20 | 1.70 | 1.80 | 0.70 |  | 5.00 |
| Paediatrics, GP referrals (1st attendance) (per 1000) | 2009/10 | 8.20 | 7.80 | 5.70 | 0.0 |  | 34.6 |
| Urology, GP Referrals (first attendance) (per 1000) | 2009/10 | 7.50 | 6.80 | 7.90 | 0.0 |  | 29.7 |
| Urology - return ratio | 2009/10 | 1.90 | 2.00 | 2.10 | 0.3 |  | 20.5 |
| A&E attendances per 1000 population (per 1000) | 2009/10 | 237.6 | 342.1 | 372.8 | 9 |  | 1191 |
| All admissions - All ages (per 1000) | 2009/10 | 161.5 | 189.1 | 231.6 | 26.5 |  | 503.0 |
| All admissions - A&E (per 1000) | 2009/10 | 8.50 | 13.2 | 9.60 | 0.0 |  | 84.5 |
| All admissions - CHD (per 1000) | 2009/10 | 4.60 | 4.40 | 4.70 | 0.0 |  | 25.9 |
| All admissions - Respiratory (per 1000) | 2009/10 | 11.8 | 13.1 | 16.0 | 1.2 |  | 51.7 |
| Elective Admission Rate (per 1000) | 2009/10 | 74.7 | 91.3 | 120.6 | 10.5 |  | 354.4 |
| Emergency Admission Rate (per 1000) | 2009/10 | 63.6 | 69.0 | 86.7 | 10.7 |  | 267.4 |
| Ambulatory Care Sensitive Conditions Rate (per 1000) | 2009/10 | 9.83 | 13.0 | 14.5 | 1.5 |  | 61.0 |
| CHD Elective Admission Rate (per 1000) | 2009/10 | | 1.80 | 1.80 | 0.00 |  | 10.00 |
| CHD Emergency Admission Rate (per 1000) | 2009/10 | 3.00 | 2.30 | 2.50 | 0.0 |  | 13.9 |
| Diabetes admission rate (per 1000) | 2009/10 | 0.000 | 0.900 | 1.10 | 0.0 |  | 10.9 |
| COPD admission rate (per 1000) | 2009/10 | | 0.900 | 2.00 | 0.0 |  | 17.2 |
| Cancer admission rate (per 1000) | 2009/10 | 18.0 | 16.7 | 28.1 | 1.3 |  | 105.2 |
| Emergency readmissions within 28d of discharge | 2009/10 | 9.1% | 8.2% | 7.7% | 0.0% |  | 31.3% |

Indicators for Needs Assessment

| Indicator | Period | Practice Value | PCT Value | England Average | England Lowest | England Range | England Highest |
|--|-----------------|----------------|-----------|-----------------|----------------|---------------|-----------------|
| Life expectancy - MSOA based estimate (Years) | 2004 - 08 | 82.5 | 83.6 | 81.8 | 75.3 | | 102.5 |
| Fertility - MSOA based estimate (per 1000) | 2003 - 07 | 55.1 | 69.1 | 59.2 | 11.1 | | 122.5 |
| Low birthweight births - MSOA based estimate | 2003 - 07 | 9.7% | 9.1% | 7.9% | 2.9% | | 13.3% |
| Smoking attributable mortality - MSOA based estimate | 2003 - 05 | 63.2 | 79.4 | 100.0 | 38.4 | | 245.9 |
| Mortality from causes amenable to healthcare - MSOA based estimate | 2003 - 05 | 92.1 | 104.4 | 100.0 | 25.6 | | 241.1 |
| Obesity prevalence - MSOA based estimate | 2003 - 05 | 15.1% | 20.7% | 23.4% | 8.7% | | 33.4% |
| Elective Admissions for All causes - MSOA based estimate | 2003/04 - 07/08 | 94.4 | 98.9 | 100.0 | 47.2 | | 168.5 |
| Emergency Admissions for All causes - MSOA based estimate | 2003/04 - 07/08 | 84.3 | 93.4 | 100.0 | 40.0 | | 219.1 |
| Elective Admissions for CHD - MSOA based estimate | 2003/04 - 07/08 | 125.5 | 115.3 | 100.0 | 36.8 | | 299.2 |
| Emergency Admissions for CHD - MSOA based estimate | 2003/04 - 07/08 | 108.0 | 138.5 | 100.0 | 41.8 | | 300.3 |
| Elective Admissions for Respiratory diseases - MSOA based estimate | 2003/04 - 07/08 | 93.1 | 91.1 | 100.0 | 29.3 | | 392.7 |
| Emergency Admissions for Respiratory diseases - MSOA based estimate | 2003/04 - 07/08 | 85.7 | 94.0 | 100.0 | 34.0 | | 293.2 |
| Emergency Admissions for Myocardial Infarction - MSOA based estimate | 2003/04 - 07/08 | 91.6 | 84.9 | 100.0 | 19.8 | | 265.9 |
| Emergency Admissions for COPD - MSOA based estimate | 2003/04 - 07/08 | 57.6 | 80.5 | 100.0 | 15.8 | | 433.1 |
| Elective Admissions for Hip replacement - MSOA based estimate | 2003/04 - 07/08 | 56.6 | 57.2 | 100.0 | 22.9 | | 198.7 |
| Emergency Admissions for Hip replacement - MSOA based estimate | 2003/04 - 07/08 | 56.4 | 56.7 | 100.0 | 37.6 | | 236.3 |
| Elective Admissions for Knee replacement - MSOA based estimate | 2003/04 - 07/08 | 127.4 | 107.8 | 100.0 | 17.2 | | 178.6 |
| Emergency Admissions for Stroke - MSOA based estimate | 2003/04 - 07/08 | 73.9 | 107.3 | 100.0 | 41.8 | | 225.5 |
| First Outpatient Attendances - MSOA based estimate | 2007/08 | 82.7 | 98.5 | 100.0 | 34.4 | | 213.0 |
| Subsequent Outpatient Attendances - MSOA based estimate | 2007/08 | 100.7 | 116.0 | 100.0 | 41.6 | | 202.4 |
| Outpatient Re-attendances - MSOA based estimate | 2007/08 | 2.67 | 2.52 | 2.24 | 1.20 | | 3.81 |